

LETTER • **OPEN ACCESS**

# Quantitative assessment of carbon sequestration reduction induced by disturbances in temperate Eurasian steppe

To cite this article: Yizhao Chen *et al* 2017 *Environ. Res. Lett.* **12** 115005

View the [article online](#) for updates and enhancements.

## Related content

- [Modeling Long-term Forest Carbon Spatiotemporal Dynamics With Historical Climate and Recent Remote Sensing Data](#)  
Jing M. Chen
- [Responses of alpine grassland on Qinghai–Tibetan plateau to climate warming and permafrost degradation: a modeling perspective](#)  
Shuhua Yi, Xiaoyun Wang, Yu Qin *et al.*
- [Disturbance-induced reduction of biomass carbon sinks of China's forests in recent years](#)  
Chunhua Zhang, Weimin Ju, Jing M Chen *et al.*

## Environmental Research Letters



## LETTER

## Quantitative assessment of carbon sequestration reduction induced by disturbances in temperate Eurasian steppe

## OPEN ACCESS

## RECEIVED

23 October 2016

## REVISED

31 July 2017

## ACCEPTED FOR PUBLICATION

7 August 2017

## PUBLISHED

31 October 2017

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Yizhao Chen<sup>1,2,3</sup>, Weimin Ju<sup>4</sup>, Pavel Groisman<sup>5</sup>, Jianlong Li<sup>2,9</sup>, Pavel Propastin<sup>6,7</sup>, Xia Xu<sup>1</sup>, Wei Zhou<sup>8</sup> and Honghua Ruan<sup>1</sup>

<sup>1</sup> Joint Innovation Center for Modern Forestry Studies, College of Biology and the Environment, Nanjing Forestry University, Nanjing, Jiangsu, People's Republic of China

<sup>2</sup> School of Life Science, Nanjing University, Nanjing, People's Republic of China

<sup>3</sup> Ministry of Education Key Laboratory for Earth System Modeling, and Department for Earth System Science, Tsinghua University, Beijing, 100084, People's Republic of China

<sup>4</sup> International Institute of Earth System Science, Nanjing University, Nanjing, People's Republic of China

<sup>5</sup> NC State University Research Scholar at National Centers for Environment Information and Hydrology Science and Services Corporation, Asheville, NC, United States of America

<sup>6</sup> Institute of Geography, Georg-August University Göttingen, Göttingen, Germany

<sup>7</sup> Department of Bioclimatology, Büsgen-Institute, Georg-August University Göttingen, Göttingen, Germany

<sup>8</sup> College of Architecture and Urban Planning, Chongqing Jiaotong University, Chongqing, People's Republic of China

<sup>9</sup> Author to whom any correspondence should be addressed.

E-mail: [jianlongli@gmail.com](mailto:jianlongli@gmail.com)

**Keywords:** temperate Eurasian steppe, carbon sequestration, disturbance simulation, terrestrial modelling, grazing activity

Supplementary material for this article is available [online](#)

**Abstract**

The temperate Eurasian steppe (TES) is a region where various environmental, social, and economic stresses converge. Multiple types of disturbance exist widely across the landscape, and heavily influence carbon cycling in this region. However, a current quantitative assessment of the impact of disturbances on carbon sequestration is largely lacking. In this study, we combined the boreal ecosystem productivity simulator (BEPS), the Shiyomi grazing model, and the global fire model (Glob-FIRM) to investigate the impact of the two major types of disturbance in the TES (i.e. domestic grazing and fire) on regional carbon sequestration. Model performance was validated using satellite data and field observations. Model outputs indicate that disturbance has a significant impact on carbon sequestration at a regional scale. The annual total carbon lost due to disturbances was  $7.8 \text{ TgCyr}^{-1}$ , accounting for 14.2% of the total net ecosystem productivity (NEP). Domestic grazing plays the dominant role in terrestrial carbon consumption, accounting for 95% of the total carbon lost from the two disturbances. Carbon losses from both disturbances significantly increased from 1999 to 2008 ( $R^2 = 0.82$ ,  $P < 0.001$  for grazing,  $R^2 = 0.51$ ,  $P < 0.05$  for fire). Heavy domestic grazing in relatively barren grasslands substantially reduced carbon sequestration, particularly in the grasslands of Turkmenistan, Uzbekistan, and the far southwest of Inner Mongolia. This spatially-explicit information has potential implications for sustainable management of carbon sequestration in the vast grassland ecosystems.

**1. Introduction**

Grassland ecosystems cover about 40% of the Earth's land surface and represent a significant carbon pool of about 20% of the world's soil carbon stocks (Wang *et al* 2008). Of these ecosystems, the Eurasian steppe belt is the largest temperate grassland eco-zone in the world and is also a wide, continuous vegetation transition zone stretching from Hungary to northeast

China. It is bounded by the forests of Siberia to the north and the deserts of Kyzylkum, Karakum, and Gobi to the south, and spans almost one-fifth of the longitudinal belt around the globe with an area of more than  $1.3 \times 10^8 \text{ km}^2$  (Wesche and Treiber 2012). This eco-zone plays an important role in the regional and global carbon cycle due to its high capacity for carbon storage and high variability in carbon source/sink conditions (Goldammer and Stocks 2000). At

present, it remains a focus of global change research (Mu *et al* 2013b, Peng *et al* 2013, Wang *et al* 2013).

The temperate Eurasian steppe (TES) is a region with various environmental, social, and economic stresses. In this vast area, disturbances induced by human beings and climate change exist widely across the landscape, and significantly affect vegetation productivity and carbon stock conditions. One of the major human activities in the grasslands is animal husbandry. According to an FAO Report (Conant 2010), grasslands are widely used as a crop and grazing source, and 27% of milk and 23% of beef produced in the world comes from grasslands. In the past 30 years, the TES has suffered from continuous degradation, partly caused by overgrazing (Conant and Paustian 2002, Gong H 2006). In some pasture ecosystems that traditionally had good forage quality and high productivity (e.g. the Xilinhot Steppe and the HulunBuir Steppe in Inner Mongolia), the quality of grass resource is now much worse than during the nomadic period (Barger *et al* 2004, He *et al* 2005). Grassland degradation and even desertification have been found in some areas of the TES during the past several decades (Mu *et al* 2013a, Zhou *et al* 2014). Obvious declines in productivity were found by field studies with medium or high stocking density of livestock (Fernandez-Gimenez and Allen-Diaz 2001, Li *et al* 2008, Li *et al* 2012). One recent study reported that domestic grazing ('grazing' from now on) reduced large amounts of vegetation carbon uptake and turned grasslands in Central Asia from carbon sinks to sources (Han *et al* 2016).

The underlying mechanisms of how grazing influences terrestrial carbon cycling stem from three processes: defoliation; carbon inputs from livestock excrement; and effects of trampling (Liu *et al* 2015, Vuichard *et al* 2007). Defoliation from grazing is the primary impact on grassland ecosystems because it directly removes biomass from plants and standing litter, which reduces carbon inputs to litter and soil carbon pools. It could also increase the carbon investment in roots as well as the CN ratio, both of which promote turnover of soil organic carbon (Mikola *et al* 2009). Excretion (urine and feces) from livestock returns carbon to the soil, and facilitates redistribution of soil nutrients (Stumpp *et al* 2005), but also negatively affects above-ground vegetation due to the urine scorching effect (Guthery and Bingham 1996). Trampling mainly changes physical and chemical properties of soil by increasing soil alkalinity, soil water holding capacity, and porosity, and thus changes the environment where microbial decomposition occurs (Francini *et al* 2014, Hiltbrunner *et al* 2012, Li *et al* 2011, Steffens *et al* 2008). Bagchi and Ritchie (2010) further proposed that changes in soil temperature, aeration, and insolation induced by grazing would indirectly influence microbial activities and decomposition. These three factors interact with each other in grasslands and determine the overall effects of grazing on the entire ecosystem.

Fire events are another potentially important disturbance that influences regional carbon sequestration. More frequent and severe steppe fires were found in Kazakhstan due to the recent warming trend (Smelansky and Tishkov 2012). This trend has been detected in the past several decades and is predicted to continue (Kamp *et al* 2016, Tchebakova *et al* 2009). In Mongolia, a multi-century chronological dataset showed that elevated temperatures did not directly increase fire events, but persistent drought events induced higher fire frequency and shorter fire return time (Hessl *et al* 2016).

Although the importance of disturbances to carbon sequestration has been recognized and supported by many field experiments, the quantitative assessment of disturbance to regional carbon sequestration is still mostly lacking (Barros and Field 2014, Li *et al* 2015, Lioubimtseva and Henebry 2009). The lack of these data contributes to uncertainty in mitigation estimates for global grasslands with a range of 0.15 to 1.3 Gt CO<sub>2</sub> yr<sup>-1</sup> (Henderson *et al* 2015, Lal 2004). In addition, the distribution of livestock has been unmanaged for many years, which has led to excessive carbon removal from the system (Lioubimtseva and Henebry 2009). Suitable manage practices are proven to be efficient for recovery and augmentation of carbon stocks, e.g. moving grazing pressure to lightly grazed areas, and growing legumes or more productive grass species (Henderson *et al* 2015, Phelan *et al* 2015). Insights into the spatial patterns of disturbance intensity and their relative importance are required to identify problem regions and properly implement sustainable management practices. High-resolution, geographically explicit mapping of livestock grazing intensity is essential to identify where mitigation efforts are urgently needed for sustainable management of grassland carbon stocks.

In order to properly understand carbon sequestration patterns in the TES and the regional impacts of two major disturbances (grazing and fire damage in this study), we combined the late version of the boreal ecosystem productivity simulator (BEPS), the Shiyomi grazing model, and the global fire model (Glob-FIRM) to quantify how carbon sequestration was changed by disturbances from 1999 to 2008. Based on the model results, we explored: (1) to what extent the two disturbances influenced carbon sequestration; and (2) the spatio-temporal pattern of disturbance intensity that may potentially be instructive for sustainable management in the region.

## 2. Materials and methods

### 2.1. Study region

The study region lays in the central part of Eurasia, spanning from Volgograd Oblast, Russia in the west to Mongolia and Inner Mongolia, China in the east (figure 1). The longitudinal and latitudinal extents are




Figure 1. Map of the study region. The green colored areas on the top map represent the grassland areas; the scale bar belongs to the top map.

39°–125°E and 35°–55°N, respectively. The region is generally dominated by a temperate continental climate with different degrees of aridity. Mean annual precipitation in much of the region is less than 400 mm. Temperature conditions have large spatiotemporal variations. Mean annual temperature ranges from  $-5^{\circ}\text{C}$  to  $15^{\circ}\text{C}$  with a latitude gradient except for the Tian Shan and Altai mountain ranges (Mohammad *et al* 2013). For seasonal variation, summer is typically hot averaging around  $20^{\circ}\text{C}$ – $26^{\circ}\text{C}$ , peak daily temperature could exceed  $40^{\circ}\text{C}$ , while in winter, the climate is extremely harsh, with low temperatures reaching  $-30^{\circ}\text{C}$ . With longitudinal differences of climate and continentality variations, zonal grassland types vary from meadow steppe (forest steppe), typical (true) steppe to desert/semi-desert steppe. Based on these longitudinal divisions of the Eurasian steppe biome, the region is separated into two sub-regions: Transvolga–Kazakhstan Steppe (TKS) and Mongol Steppe (MS) (figure 1).

## 2.2. The pixel-based livestock distribution map

Political and social factors largely influence the livestock distribution pattern at both spatial and temporal scales among different countries or sub-regions (Chen *et al* 2015b). To exclude the factors of political/social heterogeneity that control or alter the regional livestock distribution pattern, a resource-oriented scheme was developed. The basic concept is that the livestock distribution is proportional to the local grassland productivity at large scales. We first incorporated official province/prefecture level statistics to generate the regional administrative map of

livestock distribution. Thereafter, we applied a net primary productivity (NPP)-driven scheme to predict the pixel level of livestock distribution pattern.

The detailed description and validation of the livestock distribution map are presented in section S1, in the online supplementary data.

## 2.3. Model description

### 2.3.1. The BEPS model

BEPS is a remote sensing based terrestrial biogeochemical model. The model was initially developed to simulate vegetation production in boreal ecosystems (Chen *et al* 1999, Liu *et al* 1997). Then the model was coupled with terrestrial water and energy cycling and a CENTURY-derived soil carbon scheme (Chen *et al* 2007, Ju *et al* 2006, Liu *et al* 2003). It has been well validated and applied in multi-scale studies (Zhang *et al* 2012, Zhang Fangmin *et al* 2014). The model was recently revised to improve its simulation ability on grassland ecosystems (Chen *et al* 2016).

Here we only describe the algorithms to calculate the major indexes of terrestrial carbon flux used in this study.

NEP is the difference between NPP and soil respiration ( $R_h$ ):

$$\text{NEP} = \text{NPP} - R_h$$

where NPP is the difference between GPP and autotrophic respiration rate ( $R_a$ ):

$$\text{NPP} = \text{GPP} - R_a$$

GPP is calculated using the canopy ‘two-leaf’ scheme and the Farquhar biochemical model (Farquhar *et al* 1980).




Figure 2. The outline of the model pool and flux scheme in the disturbance-coupled model.

$R_a$  is simulated as the sum of growth respiration ( $R_g$ ) and maintenance respiration ( $R_m$ ).  $R_g$  is calculated as a fixed percentage from GPP. For the  $R_m$  simulation, we replaced the original scheme of  $R_m$  for grassland in BEPS with a component-specific scheme:

$$R_{m,k} = T_k \times r_{m,k} \times Q_{10,r} \frac{(T - T_{ref})}{10}.$$

$T_k$  is the biomass of the plant component  $k$ , and  $r_{m,k}$  is a coefficient of the maintenance respiration.  $T$  and  $T_{ref}$  are air temperature and reference temperature for maintenance respiration, respectively.  $Q_{10,r}$  is a temperature sensitivity factor for respiration calculated as a function of temperature (Arora 2003):

$$Q_{10,r} = 3.22 - 0.046T.$$

$R_h$  is calculated by multiplying the carbon release rate ( $p_k$ ) from the total carbon amount of soil carbon pool  $k$  ( $C_k$ ).

$$R_h = \sum_{k=1}^8 p_k C_k.$$

### 2.3.2. The grazing model

The Shiyomi grazing model (Shiyomi *et al* 2011) was modified and applied to assess the grazing consumption. This model was initially developed and successfully tested in the TES.

In the model, livestock cycling is linked to ecosystem cycling by the following processes: (1) direct consumption of aboveground biomass; (2) livestock respiration to the atmosphere; (3) excretion that adds carbon back to the soil and we further added (4) detrimental physical and chemical detrimental effects to the grassland.

Detailed algorithms of the grazing model can be found in section S2, online supplementary data available at [stacks.iop.org/ERL/12/115005/mmedia](http://stacks.iop.org/ERL/12/115005/mmedia).

### 2.3.3. The fire model

The Glob-FIRM model (Thonicke *et al* 2001) was used in this study. The model predicts the length of fire season using a conception of extinction moisture. The areas affected by fire are estimated as following:

$$N(w) = w \times e^{[k(w-1)^2]}.$$

Where  $N(w)$  is a temporal and spatial combination fraction of the fire weather for a specific pixel during the year,  $w$  is the fraction of days in a year with particular fire condition, which is calculated by the upper soil layer moisture status (Viegas *et al* 1992).

The fire effect on grasslands is calculated by the prescribed fire resistance. Above-ground biomass and litter are assumed to be fully consumed. So the carbon burned by fire can be calculated as a product of available surface carbon ( $C_{surface, ave}$ ) and  $N(w)$ :

$$C_{fire} = C_{surface, ave} \times N(w).$$

## 2.4. Model simulations

Due to quick responses and short recovery times of herbaceous plants, lagged effects of disturbances are implicitly included in the BEPS simulation (Oesterheld and McNaughton 1991, Röder *et al* 2008). In this study, we defined the impacts of disturbances as direct carbon losses due to grazing and fire. The model first simulated the grazing carbon loss and then the fire loss was then simulated after an update of the carbon pool states. At the current stage, we do not consider the interactive effect between grazing and fire events. Hence, the total disturbance impact was calculated as the sum of grazing and fire carbon consumption.

## 2.5. Pilot evaluation of modeling results

Prior to the regional scale simulations, model outputs were evaluated using several approaches. The model outputs of plant and soil carbon were validated against multi-site field observations distributed in Kazakhstan, Xinjiang, China and Inner Mongolia, China. The net ecosystem exchange (NEE, equal to minus NEP) outputs were validated using the gap-filled eddy covariance (EC) dataset.

To further test the specific performances of the two disturbance models, two other validations were set. The simulated temporal trend of grazing was validated against a multi-year fenced experiment with different livestock density levels; regional results of fire events were compared to an established remote sensing based dataset, the L3JRC product (Tansey *et al* 2008). Both the burnt area and the date of fire occurrence were validated.

## 2.6. Data use and preparation

### 2.6.1. Regional model inputs

The IGBP land cover map (Loveland *et al* 2000) was used to identify the grassland areas. The regional vegetation, meteorology, soil and livestock datasets were used to drive the model from 1999–2008, including the GlobalLAI leaf area index dataset ([www.globalmapping.org/globalLAI/](http://www.globalmapping.org/globalLAI/)), daily meteorological data from Global Meteorological Forcing Dataset for Land Surface Modeling (<http://rda.ucar.edu/datasets/ds314.0/>), monthly atmospheric CO<sub>2</sub> data from the Mauna Loa Observatory, Hawaii (20°N, 156°W) (<http://cdiac.esd.ornl.gov/ftp/trends/co2/maunaloa.co2>) and a soil texture dataset from Global Soil Dataset for use in Earth system models (<http://globalchange.bnu.edu.cn/research/soilw>). All the input data were extracted to the study region and resampled to 8 × 8 km<sup>2</sup> spatial resolution within ArcGIS 10.0 software ((ESRI) [www.esri.com/software/arcgis/](http://www.esri.com/software/arcgis/)).

### 2.6.2. Model validation data

**Flux data.** Data from nine grassland eddy covariance (EC) flux stations were collected to validate NEP results: five of them are located in the TES (in or near the study region). We also used flux data of three other stations from the temperate grassland areas. These stations share the same climate conditions with the TES according to world maps of the Köppen–Geiger climate classification (<http://koeppen-geiger.vu-wien.ac.at/>). The data were collected from the Fluxnet web site ([www.fluxdata.org/](http://www.fluxdata.org/)). We selected the flux sites with complete meteorological information needed for model simulation.

**Field research data.** 120 sites of field observations were used to validate the modeled plant and soil carbon. The sites of field measurements were distributed in eastern Kazakhstan, Xinjiang and Inner Mongolia. Plant biomass was measured in all 120 sites, and soil organic carbon was measured in 64 sites. Each

of the test sites could represent the homogenous areas of 1 hm<sup>2</sup>.

Total carbon was defined as the sum of plant and soil carbon pools. The samples were collected from four replicates for each site. The plant biomass was defined as the sum of the herbaceous, litter and root biomass. The peak-season living aboveground biomass and litter was measured by the destructive sampling of 1.0 m<sup>2</sup>. The recent litters were selected and weighted. Roots were collected by excavating to a depth of 0.5 m. Root matter was hand sorted from soil, washed, dried and weighted. The dry matter values were converted to carbon assuming a proportional factor of 0.475 (Garbulsky and Paruelo 2004).

**Fenced grazing observation.** A multi-year fenced site observation was used to evaluate the performance of the grazing model (Wang, 1998). The experiment sites located in the center of Inner Mongolia (116°04'E–117°05'E, 43°26'N–44°08'N) were under different mean stocking intensities (1.33, 2.0, 4.0, 5.33 sheep per hm<sup>2</sup>). Three independent fields of 1 hm<sup>2</sup> were set for each stocking intensity experiment. Aboveground biomass was measured using the destructive sampling method with four couples of 1 m<sup>2</sup> for each experiment field (four plots inside the field and four plots outside the field). The observations were taken in 1990, 1993 and 1997. The experiment started May and ended on October in each year, and biomass was measured every 15 days.

**Fire regime dataset.** The L3JRC product (Tansey *et al* 2008) was used to validate the performance of the fire model. The dataset is based on SPOT VEGETATION Earth Observation system with 1 km<sup>2</sup> spatial resolution and daily intervals. The product also records the date of fire occurrence. The temporal extension is from 2000–2007.


All sites' information is shown in table S2 in the online supplementary data.

## 3. Results

### 3.1. Model validations

Comparisons between modeled and observed results are shown in figure 3. The modeled results were consistent with field measurements. The  $R^2$  values are 0.84 and 0.58 for plant and soil C, respectively. The corresponding slopes are 0.74 and 0.78.

Model outputs of NEE were generally consistent with continuous observations at the eight grassland flux sites (figure 4), and we were able to simulate temporal tendencies in the model. Modeled annual net ecosystem exchange ( $NEE_{MOD}$ ) values were higher than measured values ( $NEE_{EC}$ ). The  $R^2$  values between  $NEE_{MOD}$  and  $NEE_{EC}$  ranged from 0.616 (US\_BKG) to 0.848 (RU\_HA2), and RMSE values ranged from 0.4 gC m<sup>-2</sup> d<sup>-1</sup> (RU\_HA1) to 1.307 gC m<sup>-2</sup> d<sup>-1</sup> (US\_BKG) (figure 4(b)).



Comparisons against the fenced observations show that our model set could reproduce seasonal trends of grassland production under different levels of stocking pressure (figure 5). The overall  $R^2$  is 0.71 ( $P < 0.01$ ).

Comparisons of burned areas between the L3JRC dataset and model simulation are shown in (figure 6 (a)); 86% of the pixels are in agreement with the L3JRC dataset. We also compared the regional mean fire occurrence day from 2000–2007; the model identified the regional temporal pattern when fire starts to appear ( $R^2 = 0.52$ ,  $P < 0.05$ , figure 6(b)).

### 3.2. Regional pattern of NPP and NEP

Spatial distributions of mean NPP and NEP from 1999–2008 are shown in figure 7. As a widespread biome transition belt, the TES connects deserts in the southwest and forests in the north; vegetation productivity varies based on these regions. High productivity grasslands are mainly concentrated in areas that are connected with forested areas, such as: meadow steppe in the eastern part of the Mongol Steppe, which is joined to Great Khingan and Siberia; alpine meadow steppe around the Tian Shan and Altai Mountains; and Russian steppe areas that are close to the North Caucasus Forest. Low productivity areas are distributed around the deserts of Kyzylkum and Karakum in the TKS and the Gobi Desert in MS (figure 7(a1)). The model-calculated spatial pattern of NEP is basically consistent with NPP, but may be modified in part by the heterotrophic respiration distribution (figure 7(b1)). Low productivity areas generally acted as carbon sources or were neutral, and high productivity areas acted as carbon sinks. The major differences of spatial distribution were located in the northern provinces in Mongolia (Dornod and Hentiy) and Kazakhstan (Pavlodar and Qaraghandy). High productivity did not result in higher carbon sequestration in the southwest part of the Russian

grasslands due to the high soil respiration rate in the red soil of the North Caucasus (Alexandrovskiy *et al* 2014).


Based on these results, the mean and total NPP of TES were  $88.41 \text{ gC m}^{-2} \text{ yr}^{-1}$  and  $392 \text{ TgC yr}^{-1}$  for the study period. The mean NPP in the TKS and the MS were similar:  $87.7 \text{ gC m}^{-2} \text{ yr}^{-1}$  and  $89.5 \text{ gC m}^{-2} \text{ yr}^{-1}$ , respectively. The total NPP in these two regions was  $240 \text{ TgC yr}^{-1}$  and  $152 \text{ TgC yr}^{-1}$ , respectively (figure (a2)).

The decadal mean NEP of TES was  $12.86 \text{ gC m}^{-2} \text{ yr}^{-1}$ , indicating that the region was a net carbon sink. The total NEP was  $39.2 \text{ TgC yr}^{-1}$ . The values were positive for both sub-regions. The total NEP for the TKS and the MS was  $14.4 \text{ TgC yr}^{-1}$  and  $24.8 \text{ TgC yr}^{-1}$ , respectively (figure 7).

### 3.3. The regional patterns of grazing and fire disturbances and their impacts on carbon sequestration

The spatial distribution of the two types of disturbances was very uneven in the TES. For the fire regime, the major influenced areas were the northern and north-eastern TKS. However in the MS, fire rarely happened (figure 7(a3)). For grazing, there was a distinct spatial imbalance in carbon consumption in different countries and in different provinces or prefectures (figure 7(a4)). In the MS, major consumption was concentrated in China, especially in Inner Mongolia. The most extensively used grasslands were found in the traditional natural pastures with relatively high productivity, such as XilinGol Prairie in mid-Inner Mongolia and Hulunbuir Prairie close to the Great Khingan. In the TKS, high consumption was found in Uzbekistan and Turkmenistan, where two deserts (Kyzylkum and Karakum) are located and the major grassland type is desert steppe.

Both types of disturbance increased significantly during the study period ( $R^2 = 0.82$ ,  $P < 0.001$  for grazing,  $R^2 = 0.51$ ,  $P < 0.05$  for fire, figure 8). For





grazing, carbon loss was on a near-linear increasing trend in the TKS; and for the MS, the trend was similar, except for a curved valley in 2002, which was probably caused by a long-term dzud (severe winter) in Mongolia in that year (Tachiiri *et al* 2008). For the fire regime, the main affected region was the TKS, and NPP loss from fire increased significantly over the decade of the study. In the TKS, the means and total disturbance consumption values were  $1.76 \text{ gC m}^{-2}$

$\text{yr}^{-1}$  and  $7.8 \text{ TgCyr}^{-1}$ , respectively (figure 7(b3)). Carbon consumption from grazing was much higher than for fire, and accounted for 95% of the total carbon loss.

For the first seven years, the region acted as a carbon sink, but in the last three years it turned into a weak source. This trend was mainly caused by a rapid NEP decrease, from  $34.56 \text{ gC m}^{-2}$  to  $-30.96 \text{ gC m}^{-2}$  in the TKS during between 2002–2008 (with a slight





increase in 2007). The inter-annual results indicated that the disturbances alleviated the regional carbon sink from 1999–2005 and aggravated the carbon source effect from 2006–2008 (figure 9).

The disturbance impact was much larger in the MS than in the TKS due to the higher level of grazing in the MS sub-region. According to our results, the mean percentage of grazing consumption from the NEP was 13.5% during the decade. The values were 7.6% and 16.6% for the TKS and the MS, respectively. For fire events, the percentages were 0.7% for the entire region, 0.86% for the TKS, and 0.2% for the MS. If the carbon losses from disturbances were explicitly considered, regional carbon sequestration decreased from  $39.2 \text{ Tg C yr}^{-1}$  to  $31.4 \text{ Tg C yr}^{-1}$ .

The spatial distribution of the impact of grazing to NEP is displayed in figure 10. Grazing could reverse the local carbon sink to a source in some areas, if the grazing consumption was higher than the absolute value of NEP. This phenomenon is especially significant in barren areas, such as the grasslands in Turkmenistan and Uzbekistan and the southwest boarder of grasslands in Inner Mongolia, China.

We further quantified the grazing impact to NEP for each country and sub-region and found a great spatial imbalance presented in figure 10. Grazing has minor impacts in Kazakhstan, Kyrgyzstan, Russia, Xinjiang, China and Mongolia with a percentage lower than 1% from NEP. However, in arid countries like Uzbekistan and Turkmenistan, over-grazing widely exists. Grazing consumption exceeded NEP in 71.9% and 92.6% of grasslands in these two countries, respectively. These results demonstrate the strong impact of grazing in altering the carbon budget in these areas.

## 4. Discussions

### 4.1. Recent trends and characteristics of disturbances in the TES

The two types of disturbances we studied lead to profound impacts on carbon sequestration in the TES. If the carbon losses from disturbances were explicitly considered, carbon sequestration was significantly reduced in this region. According to our study, the




Figure 7. Spatial distribution of mean annual (a) NPP, (b) NEP, (c) grazing consumption and (d) fire consumption.




Figure 8. Variation in annual total carbon consumption due to grazing and fire from 1999–2008.




Figure 9. The impact of disturbances to regional NEP of (a) the TES, (b) the TKS, and (c) the MS from 1999–2008.

total consumption from the two types of disturbances was  $7.8 \text{ Tg C yr}^{-1}$ , accounting for 14.2% of the regional NEP, and grazing is the major contributor to carbon loss. Although the importance of grazing has been recognized for a long time, it has not been well quantified into the regional or global carbon sequestration calculations. Therefore, our results could contribute to understanding the large uncertainty in carbon sequestration abilities in arid and semi-arid ecosystems around the world (Ahlström *et al* 2015).

Chen *et al* (2015b) proposed the concept of coupled human and natural systems (CHANS), which focuses on the strong interactions between biophysical and socioeconomic systems among TES regions. The CHANS framework suggests that the importance of anthropogenic influences might produce stronger impacts on the grasslands than climate change (Chen *et al* 2015a). Although it is still difficult to explicitly disentangle the relative contributions of the two factors, our study provides strong quantitative evidence for the anthropogenic part.




Figure 10. Map (a) and statistical summary (b) of the percentage of grazing consumption from the absolute values of NEP.

Furthermore, the carbon consumption caused by both types of disturbance significantly increased during the study period. These trends correspond to the official statistics and previous observations. FAO statistics (FAOSTAT, <http://faostat.fao.org/>) showed that the livestock numbers in all nations and regions in TES have increased since the late 1990s. Chen *et al* (2013) reported that more than 40% of grasslands in Central Asia were used for animal husbandry in 2009 compared to 2000 based on remote sensing data. With the continually expanding demand of livestock products and a 'warming and drying' trend, both the scale of animal husbandry and fire danger are predicted to increase (Flannigan *et al* 2013, Godber and Wall 2014). Hence, the effects of disturbance in TES will probably continue to expand and strongly impact regional carbon cycling in the future.

Moreover, the uneven distribution of disturbances and different management styles need to be considered and mitigated. Our results suggested that similar net grazing consumption occurred in southern countries in the TKS and in Inner Mongolia, China in the MS, but the effect of grazing on carbon sequestration in the southern TKS was much higher than in Inner Mongolia. These regional patterns are related to

fundamental differences in grassland quality and productivity. But more importantly, the absolute number of livestock in the two countries (i.e. Uzbekistan and Turkmenistan) is far beyond a suitable stocking rate, leading to the overall over-grazing condition. We have not found detailed statistics from these two countries, however, it is reported that the average stocking rate near settlements in Kyrgyzstan is about 2.74 sheep head  $\text{hm}^{-2}$ , which is 3.2 times higher than the recommended scale (0.85 sheep head  $\text{hm}^{-2}$ ) (Mirzabaev *et al* 2016). This result may reflect uncontrolled and excessive stocking rates in Uzbekistan and Turkmenistan, since livestock density in the two countries is obviously higher than in Kyrgyzstan. Because the grasslands are managed by state-controlled cooperatives or farmer associations, this situation is closely related to the national condition. Historically, the national economy in the two countries largely has relied on grassland resources since the collapse of the USSR (Lioubimtseva and Henebry 2009). The huge numbers of livestock and years of over-utilization has directly caused the deserts to expand (Robinson *et al* 2003). Now grassland resources are diminished but livestock numbers continue to increase after the USSR has collapsed (Chen *et al* 2013).

**Table 1.** Recent model results of grassland in the Eurasian steppe and comparison with this study.

Study area	Study period	Method	Model type of carbon assimilation <sup>a</sup>	Results (gC m <sup>-2</sup> yr <sup>-1</sup> )	Corresponding results in this study (gC m <sup>-2</sup> yr <sup>-1</sup> ) <sup>b</sup>	Reference
Mongolia	2000–2005	Improved CASA model	LUE	NPP: 125.33	75.2	(Xing <i>et al</i> 2010)
Northern China	2000–2005	Improved CASA model	LUE	NPP:153.26	113.8	(Gao and Liu 2008)
		CASA	LUE	NPP:245	121.23 (for Inner Mongolia)	
China	2007	CEVSA	Farquhar	NPP:208	121.23 (for Inner Mongolia)	(Feng <i>et al</i> 2007)
		GLOPEM	Farquhar	NPP:145		
		GEOLUE	LUE	NPP:178		
		GEOPRO	Farquhar	NPP:168		
Eastern Kazakhstan	2004	Modified light use efficiency (LUE) model	LUE	NPP:168	152.1	(Propastin <i>et al</i> 2012)
Xilinhot, Inner Mongolia	2002	GLPM	Farquhar	NEP:1.91	2.23	(Zhang <i>et al</i> 2009)
Inner Mongolia	1951–2007	TEM model	Farquhar	NEP:11.25	15.97 for the period of 1999 to 2007	(Sui <i>et al</i> 2013)

<sup>a</sup>LUE: light use efficiency model; Farquhar: Farquhar biochemical model.

<sup>b</sup>The results are extracted from the simulated map in this study with the same spatiotemporal scale to the corresponding recent studies.

Pasture management practices are the other important contributor to the current regional patterns. In China, livestock numbers have experienced a boom period since the 2000s to meet the social and economic needs for animal products in China (Rae and Zhang 2009). To alleviate pressure from unmanaged pasture use and to preserve grassland resources, the Chinese government has promoted several regional programs like the Grain to Green Program (GTGP) and the Grazing Withdrawal Program (GWP) since the 1990s. The primary goal of both programs is to conserve pastures under a different level of degradation by transforming grazing and cropping activity to be sustainable. According to our previous study in Inner Mongolia, construction of infrastructure, including animal sheds and corrals, increased by 20%, and the size of fenced grassland increased by nearly three times from 1999–2008. Accordingly, grassland NPP increased by 40.6 gC m<sup>-2</sup> yr<sup>-1</sup> between 2001 and 2009 (with climatic factors excluded) (Mu *et al* 2013c). On the other side of TES, in the TKS, the influence of high grazing intensity on carbon cycling has not received enough attention from the scientific community or from government officials. Although some management suggestions to conserve carbon stocks, such as rotational and migratory grazing, have been supported by empirical evidence (ICARDA 2001, 2002), field studies are still urgently needed in arid areas. Furthermore, the huge amount of investment needed to carry out rehabilitation might be a greater obstacle to reversing degradation on a large scale (Blench and Sommer 1999, Mirzabaev *et al* 2016). Over the long term, economic development is the primary task for developing countries in this region, but finding a

balance between living improvements of human beings and ecosystem sustainability is an essential topic to consider.

#### 4.2. Model uncertainty and potential improvements

We validated model outputs in several ways. The results showed general consistency of our model with continuous carbon flux observations, field data, and remote sensing product. However, model uncertainties are sometimes unavoidable when simulating a complex system with multiple factors. First, uncertainty exists in the model input of leaf area index (LAI). Unstable surface reflectance, foliage clumping, and soil effects in regions with sparse vegetation are among the major sources of LAI deviation (Liu *et al* 2012). The latter two factors generally cause underestimation of LAI, which could have caused the underestimations of NPP and NEP in this study. Second, the uncertainty could arise from mathematical descriptions and corresponding parameterization of biogeochemical processes in the model. We compared our model results with previous studies in this region (table 1). Our results are more comparable with models that share the same photosynthesis scheme (i.e. Farquhar model), but our results are often lower than light use efficiency (LUE) models.

In the current soil carbon model, the mechanical description of microbial activities is relatively weak in soil carbon circulation and the algorithms to calculate environmental factors on soil decomposition are not well supported (Luo *et al* 2015). Moreover, the disturbance intensity, especially the stocking rate, could significantly affect the magnitude of changes in carbon pools and fluxes. Light grazing might contribute to accumulation of carbon, whereas

moderate and heavy grazing would induce carbon loss in both plant and soil pools and lead to soil erosion and degradation (Yan *et al* 2013, Zhou *et al* 2016). Responses of grasslands to different disturbance intensities should be considered in future models and simulations.

With the input of data based on governmental official statistics, regional livestock consumption could be constrained at a certain range. The model validation results also support the effectiveness of the grazing model. However, the current model assumes a constant daily intake rate of livestock, which ignores the effects of environmental change such as climatic warming and CO<sub>2</sub> enhancement on the livestock intake rate (Collier and Zimelman 2007, Nardone *et al* 2010). In addition, more frequent extreme weather events cause more severe stresses that are closely related to higher livestock mortality (Vitali *et al* 2009).

For the fire regime, the major disagreement between the model results and the field observations is from the mountainous areas in north Kazakhstan and Kyrgyzstan. We suspect that deviations in the model come from the discrepancy between the model scheme and real climate patterns in these areas. The model used soil water content as a scalar to determine fire occurrence. That is, if surface soil water content is lower than 'extinction moisture', then the area has the potential to burn. In these mountainous areas, the climate pattern is different from the rest of the region. Rainfall mainly happens in the winter and spring, and the temperature is mild throughout the year, rarely exceeding 30 °C, even during the mid-summer (Rachkovskaya and Bragina 2012). Thus it is possible that the soil water content is low but energy conditions for fire occurrence are not met.

The period of our study avoids significant land use and cover change (LUCC) from political events, e.g. the collapse of the former USSR. However, some studies have highlighted the potential impacts from LUCC in recent years. For example, Chen *et al* (2013) reported that the grasslands were gradually reclaimed after the overall recovery during the 1990s in Central Asia. Mu *et al* (2013c) suggested that the increasing grassland areas lead to carbon accumulation in Inner Mongolia. Based on our results, we speculate that persistent over-grazing will largely accelerate irreversible land degradation, especially in the southern countries in the TKS. In addition, the emergence of mining and oil production, as well as the current trend of urbanization are also potential factors affecting LUCC in the TES (Peng *et al* 2009, Pomfret 2016, Suzuki 2013).

## 5. Conclusions and perspectives

In this study, we found that grassland carbon sequestration could be significantly reduced in the TES if carbon losses from major disturbance events

(grazing and fire) were explicitly considered. Grazing dominated the carbon losses from the two disturbances with a percentage of 95%. Although uncertainties exist in the model, the huge impacts of disturbances are quantified. Based on the literature search result from the ISI Web of Knowledge database ([www.isiwebofknowledge.com](http://www.isiwebofknowledge.com)), this study is among the first regional quantitative assessment of the disturbance impact to carbon sequestration in the TES. We pinpointed major over-grazing areas in the TES, which is a first step towards achieving sustainable management of regional grasslands. A potential further study would apply different grazing scenarios to see how management practices contribute to conservation of terrestrial carbon stocks.

As summarized by Qi *et al* (2012), regional grassland ecosystems are especially vulnerable to both climate change and intensified human activities. Our results support the idea that disturbance has the potential to change terrestrial ecosystems from carbon sinks to carbon sources in some areas. At the same time, our assessment of disturbance is not completely inclusive. For example, grazing mainly represents the disturbance from animal husbandry during the growing season, but during the non-growing season, livestock mainly rely on drylot feed. Total carbon loss should be carefully considered, because forage could be locally provided or imported from outside the region. Other disturbances due to mining and the petroleum industry have gradually been established in the TES since the late 1990s. The fundamental ways that mining and the petroleum industry influence grasslands could be very different from the disturbances we focused on in this study. These potential differences need to be considered in future studies.

## Acknowledgments

This work is supported by the National Key R&D Program of China (2016YFA0600202), the Fundamental Research Funds for the Central Universities (KYZ201758) the National Key R&D Program of China (2016YFD0600204), APN Global Change Fund Project (No. ARCP2013-16NMY-Li), National Youth Science Fund of China (41501575) and the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD).

We acknowledge all producers and providers of the datasets used in this study. We also greatly acknowledge Dr Jingming Chen from the University of Toronto for offering the model codes of BEPS and useful instructions, Dr Yiqi Luo from the University of Oklahoma and Dr Jianyang Xia from the East China Normal University who gave suggestions. The model design and construction is largely inspired by the topics and discussions during the NEESPI/NEFI events. Without this kind help and encouragement, this work could not be finished.

## References

- Ahlström A, Raupach M R, Schurgers G, Smith B, Arneth A, Jung M, Reichstein M, Canadell J G, Friedlingstein P and Jain A K 2015 The dominant role of semi-arid ecosystems in the trend and variability of the land CO<sub>2</sub> sink *Science* **348** 895–9
- Alexandrovskiy A, Sedov S and Shishkov V 2014 The development of deep soil processes in ancient kurgans of the North Caucasus *CATENA* **112** 65–71
- Arora V K 2003 Simulating energy and carbon fluxes over winter wheat using coupled land surface and terrestrial ecosystem models *Agric. Forest Meteorol.* **118** 21–47
- Bagchi S and Ritchie M E 2010 Herbivore effects on above- and belowground plant production and soil nitrogen availability in the trans-Himalayan shrub-steppes *Oecologia* **164** 1075–82
- Barger N N, Ojima D S, Belnap J, Shiping W, Yanfen W and Chen Z 2004 Changes in plant functional groups, litter quality, and soil carbon and nitrogen mineralization with sheep grazing in an Inner Mongolian grassland *J. Range Manage.* **57** 613–9
- Barros V and Field C 2014 *Climate Change 2014: Impacts, Adaptation, and Vulnerability: Working Group II Contribution to the IPCC 5th Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge: Cambridge University Press)
- Blench R and Sommer F 1999 *Understanding Rangeland Biodiversity* (London: Overseas Development Institute)
- Chen B, Chen J M, Mo G, Yuen C-W, Margolis H, Higuchi K and Chan D 2007 Modeling and scaling coupled energy, water, and carbon fluxes based on remote sensing: an application to Canada's landmass *J. Hydrometeorol.* **8** 123–43
- Chen J, John R, Shao C, Fan Y, Zhang Y, Amarjargal A, Brown D G, Qi J, Han J and Laforteza R 2015a Policy shifts influence the functional changes of the CNH systems on the Mongolian plateau *Environ. Res. Lett.* **10** 085003
- Chen J *et al* 2015b Divergences of two coupled human and natural systems on the Mongolian plateau *Bioscience* **65** 559–70
- Chen J, Liu J, Cihlar J and Goulden M 1999 Daily canopy photosynthesis model through temporal and spatial scaling for remote sensing applications *Ecol. Model.* **124** 99–119
- Chen X, Bai J, Li X, Luo G, Li J and Li B L 2013 Changes in land use/land cover and ecosystem services in Central Asia during 1990–2009 *Curr. Opin. Environ. Sustain.* **5** 116–27
- Chen Y, Mu S, Sun Z, Gang C, Li J, Padarian J, Groisman P, Chen J and Li S 2016 Grassland carbon sequestration ability in China: a new perspective from terrestrial aridity zones *Rangeland Ecol. Manage.* **69** 84–94
- Collier R J and Zimelman R B 2007 Heat stress effects on cattle: What we know and what we don't know *22nd Annual Southwest Nutrition & Management Conference* 76–83
- Conant R T 2010 Challenges and opportunities for carbon sequestration in grassland systems *Food and Agriculture Organization of the United Nations*
- Conant R T and Paustian K 2002 Potential soil carbon sequestration in overgrazed grassland ecosystems *Glob. Biogeochem. Cycles* **16** 9091–99
- Farquhar G, von Caemmerer S V and Berry J 1980 A biochemical model of photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species *Planta* **149** 78–90
- Feng X, Liu G, Chen J, Chen M, Liu J, Ju W, Sun R and Zhou W 2007 Net primary productivity of China's terrestrial ecosystems from a process model driven by remote sensing *J. Environ. Manage.* **85** 563–73
- Fernandez-Gimenez M and Allen-Diaz B 2001 Vegetation change along gradients from water sources in three grazed Mongolian ecosystems *Plant Ecol.* **157** 101–18
- Flannigan M, Cantin A S, de Groot W J, Wotton M, Newbery A and Gowman L M 2013 Global wildland fire season severity in the 21st century *Forest Ecol. Manage.* **294** 54–61
- Francini G, Liiri M, Männistö M, Stark S and Kytöviita M-M 2014 Response to reindeer grazing removal depends on soil characteristics in low Arctic meadows *Appl. Soil Ecol.* **76** 14–25
- Gao Z and Liu J 2008 Simulation study of China's net primary production *Chin. Sci. Bull.* **53** 434–43
- Garbulsky M F and Paruelo J M 2004 Remote sensing of protected areas to derive baseline vegetation functioning characteristics *J. Veg. Sci.* **15** 711–20
- Godber O F and Wall R 2014 Livestock and food security: vulnerability to population growth and climate change *Glob. Change Biol.* **20** 3092–102
- Goldammer J G and Stocks B J 2000 Eurasian perspective of fire: dimension, management, policies, and scientific requirements *Fire, Climate Change, and Carbon Cycling in the Boreal Forest* (Berlin: Springer) pp 49–64
- Gong H and Wang D 2006 Perspectives in optimization models of grassland grazing systems *Acta Prataculturae Sinica* **15** 1–8
- Guthery F S and Bingham R L 1996 A theoretical basis for study and management of trampling by cattle *J. Range Manage.* **49** 264–9
- Han Q, Luo G, Li C, Shakir A, Wu M and Saidov A 2016 Simulated grazing effects on carbon emission in Central Asia *Agric. Forest Meteorol.* **216** 203–14
- He C, Zhang Q, Li Y, Li X and Shi P 2005 Zoning grassland protection area using remote sensing and cellular automata modeling—a case study in Xilingol steppe grassland in northern China *J. Arid. Environ.* **63** 814–26
- Henderson B B, Gerber P J, Hilinski T E, Falcucci A, Ojima D S, Salvatore M and Conant R T 2015 Greenhouse gas mitigation potential of the world's grazing lands: modeling soil carbon and nitrogen fluxes of mitigation practices *Agric. Ecosyst. Environ.* **207** 91–100
- Hessl A E, Brown P, Byambasuren O, Cockrell S, Leland C, Cook E, Nachin B, Pederson N, Saladyga T and Suran B 2016 Fire and climate in Mongolia (1532–2010 Common Era) *Geophys. Res. Lett.* **43** 6519–27
- Hiltbrunner D, Schulze S, Hagedorn F, Schmidt M W and Zimmermann S 2012 Cattle trampling alters soil properties and changes soil microbial communities in a Swiss sub-alpine pasture *Geoderma* **170** 369–77
- ICARDA 2001 Integrated feed and livestock production in the steppes of Central Asia *IFAD Technical Assistance Grant: ICARDA-425 Annual Report (2000–2001)* 1–161
- ICARDA 2002 Integrated feed and livestock production in the steppes of Central Asia *IFAD Technical Assistance Grant: ICARDA-425 Annual Report (2001–2002)* 1–207
- Ju W, Chen J M, Black T A, Barr A G, Liu J and Chen B 2006 Modelling multi-year coupled carbon and water fluxes in a boreal aspen forest *Agric. Forest Meteorol.* **140** 136–51
- Kamp J, Koshkin M A, Bragina T M, Katzner T E, Milner-Gulland E, Schreiber D, Sheldon R, Shmalenko A, Smelansky I and Terraube J 2016 Persistent and novel threats to the biodiversity of Kazakhstan's steppes and semi-deserts *Biodivers. Conserv.* **1–21**
- Lal R 2004 Soil carbon sequestration impacts on global climate change and food security *Science* **304** 1623–7
- Li C, Hao X, Zhao M, Han G and Willms W D 2008 Influence of historic sheep grazing on vegetation and soil properties of a desert steppe in Inner Mongolia *Agric. Ecosyst. Environ.* **128** 109–16
- Li C, Zhang C, Luo G, Chen X, Maisupova B, Madaminov A A, Han Q and Djenbaev B M 2015 Carbon stock and its responses to climate change in Central Asia *Glob. Change Biol.* **21** 1951–67
- Li S, Verburg P H, Lv S, Wu J and Li X 2012 Spatial analysis of the driving factors of grassland degradation under conditions of climate change and intensive use in Inner Mongolia, China *Reg. Environ. Change* **12** 461–74

- Li Y, Zhao H, Zhao X, Zhang T, Li Y and Cui J 2011 Effects of grazing and livestock exclusion on soil physical and chemical properties in desertified sandy grassland, Inner Mongolia, northern China *Environ. Earth Sci.* **63** 771–83
- Lioubimtseva E and Henebry G M 2009 Climate and environmental change in arid Central Asia: impacts, vulnerability, and adaptations *J. Arid. Environ.* **73** 963–77
- Liu J, Chen J, Cihlar J and Park W 1997 A process-based boreal ecosystem productivity simulator using remote sensing inputs *Remote Sens. Environ.* **62** 158–75
- Liu J, Chen J M and Cihlar J 2003 Mapping evapotranspiration based on remote sensing: an application to Canada's landmass *Water Resour. Res.* **39** 1189
- Liu N, Kan H M, Yang G W and Zhang Y J 2015 Changes in plant, soil, and microbes in a typical steppe from simulated grazing: explaining potential change in soil C *Ecol. Monogr.* **85** 269–86
- Liu Y, Liu R and Chen J M 2012 Retrospective retrieval of long-term consistent global leaf area index (1981–2011) from combined AVHRR and MODIS data *J. Geophys. Res.: Biogeosci.* **117** G04003
- Loveland T, Reed B, Brown J, Ohlen D, Zhu Z, Yang L and Merchant J 2000 Development of a global land cover characteristics database and IGBP DISCover from 1 km AVHRR data *Int. J. Remote. Sens.* **21** 1303–30
- Luo Y, Keenan T F and Smith M 2015 Predictability of the terrestrial carbon cycle *Glob. Change Biol.* **21** 1737–51
- Mikola J, Setälä H, Virkajärvi P, Saarijärvi K, Ilmarinen K, Voigt W and Vestberg M 2009 Defoliation and patchy nutrient return drive grazing effects on plant and soil properties in a dairy cow pasture *Ecol. Monogr.* **79** 221–44
- Mirzabaev A, Ahmed M, Werner J, Pender J and Louhaichi M 2016 Rangelands of Central Asia: challenges and opportunities *J. Arid Land* **8** 93–108
- Mohammad A, Wang X, Xu X, Peng L, Yang Y, Zhang X, Myneni R B and Piao S 2013 Drought and spring cooling induced recent decrease in vegetation growth in Inner Asia *Agric. Forest Meteorol.* **178** 21–30
- Mu S, Yang H, Li J, Chen Y, Gang C, Zhou W and Ju W 2013a Spatio-temporal dynamics of vegetation coverage and its relationship with climate factors in Inner Mongolia, China *J. Geog. Sci.* **23** 231–46
- Mu S J, Chen Y Z, Li J L, Ju W M, Odeh I O A and Zou X L 2013b Grassland dynamics in response to climate change and human activities in Inner Mongolia, China between 1985 and 2009 *Rangeland J.* **35** 315–29
- Mu S J, Zhou S X, Chen Y Z, Li J L, Ju W M and Odeh I O A 2013c Assessing the impact of restoration-induced land conversion and management alternatives on net primary productivity in Inner Mongolian grassland, China *Glob. Planet. Change* **108** 29–41
- Nardone A, Ronchi B, Lacetera N, Ranieri M S and Bernabucci U 2010 Effects of climate changes on animal production and sustainability of livestock systems *Livest. Sci.* **130** 57–69
- Oosterheld M and McNaughton S 1991 Effect of stress and time for recovery on the amount of compensatory growth after grazing *Oecologia* **85** 305–13
- Peng S, Piao S, Shen Z, Ciais P, Sun Z, Chen S, Bacour C, Peylin P and Chen A 2013 Precipitation amount, seasonality and frequency regulate carbon cycling of a semi-arid grassland ecosystem in Inner Mongolia, China: a modeling analysis *Agric. Forest Meteorol.* **178–179** 46–55
- Peng S, Piao S, Wang T, Sun J and Shen Z 2009 Temperature sensitivity of soil respiration in different ecosystems in China *Soil Biol. Biochem.* **41** 1008–14
- Phelan P, Moloney A, McGeough E, Humphreys J, Bertilsson J, O'Riordan E and O'Kiely P 2015 Forage legumes for grazing and conserving in ruminant production systems *Crit. Rev. Plant. Sci.* **34** 281–326
- Pomfret R W 2006 *The Central Asian Economies Since Independence* (Princeton, NJ: Princeton University Press)
- Propastin P, Kappas M W, Herrmann S M and Tucker C J 2012 Modified light use efficiency model for assessment of carbon sequestration in grasslands of Kazakhstan: combining ground biomass data and remote-sensing *Int. J. Remote Sens.* **33** 1465–87
- Qi J, Chen J, Wan S and Ai L 2012 Understanding the coupled natural and human systems in dryland East Asia *Environ. Res. Lett.* **7** 015202
- Röder A, Hill J, Duguy B, Alloza J A and Vallejo R 2008 Using long time series of Landsat data to monitor fire events and post-fire dynamics and identify driving factors: a case study in the Ayora region (Eastern Spain) *Remote Sens. Environ.* **112** 259–73
- Rachkovskaya E I and Bragina T M 2012 Steppes of Kazakhstan: diversity and present state *Eurasian Steppes: Ecological Problems and Livelihoods in a Changing World* (Dordrecht: Springer Netherlands) pp 103–48
- Rae A N and Zhang X 2009 China's booming livestock industry: household income, specialization, and exit *Agric. Econ.* **40** 603–16
- Robinson S, Milner-Gulland E J and Alimaev I 2003 Rangeland degradation in Kazakhstan during the Soviet era: re-examining the evidence *J. Arid. Environ.* **53** 419–39
- Shiyomi M, Akiyama T, Wang S, Yiruhan A, Hori Y, Chen Z, Yasuda T, Kawamura K and Yamamura Y 2011 A grassland ecosystem model of the Xilingol steppe, Inner Mongolia, China *Ecol. Model.* **222** 2073–83
- Smelansky I E and Tishkov A A 2012 The steppe biome in Russia: ecosystem services, conservation status, and actual challenges, Eurasian steppes *Ecological Problems and Livelihoods in a Changing World* (Berlin: Springer) pp 45–101
- Steffens M, Kölbl A, Totsche K U and Kögel-Knabner I 2008 Grazing effects on soil chemical and physical properties in a semiarid steppe of Inner Mongolia (PR China) *Geoderma* **143** 63–72
- Stumpp M, Wesche K, Retzer V and Miede G 2005 Impact of grazing livestock and distance from water source on soil fertility in southern Mongolia *Mount. Res. Dev.* **25** 244–51
- Sui X, Zhou G and Zhuang Q 2013 Sensitivity of carbon budget to historical climate variability and atmospheric CO<sub>2</sub> concentration in temperate grassland ecosystems in China *Clim. Change* **117** 259–72
- Suzuki Y 2013 Conflict between mining development and nomadism in Mongolia *The Mongolian Ecosystem Network* (Berlin: Springer) pp 269–294
- Tachiiri K, Shinoda M, Klinkenberg B and Morinaga Y 2008 Assessing Mongolian snow disaster risk using livestock and satellite data *J. Arid. Environ.* **72** 2251–63
- Tansev K, Grégoire J M, Defourny P, Leigh R, Pekel J F, Van Bogaert E and Bartholomé E 2008 A new, global, multi-annual 2000–7 burnt area product at 1 km resolution *Geophys. Res. Lett.* **35** L01401
- Tchebakova N, Parfenova E and Soja A 2009 The effects of climate, permafrost and fire on vegetation change in Siberia in a changing climate *Environ. Res. Lett.* **4** 045013
- Thonicke K, Venevsky S, Sitch S and Cramer W 2001 The role of fire disturbance for global vegetation dynamics: coupling fire into a dynamic global vegetation model *Glob. Ecol. Biogeogr.* **10** 661–77
- Viegas D, Viegas M and Ferreira A 1992 Moisture content of fine forest fuels and fire occurrence in central Portugal *Int. J. Wildland Fire* **2** 69–86
- Vitali A, Segnalini M, Bertocchi L, Bernabucci U, Nardone A and Lacetera N 2009 Seasonal pattern of mortality and relationships between mortality and temperature-humidity index in dairy cows *J. Dairy Sci.* **92** 3781–90
- Vuichard N, Soussana J F O, Ciais P, Viovy N, Ammann C, Calanca P, Clifton Brown J, Fuhrer J R, Jones M and Martin C C 2007 Estimating the greenhouse gas fluxes of European grasslands with a process-based model: 1. Model evaluation from in situ measurements *Glob. Biogeochem. Cycles* **21**
- Wang J, Brown D G and Chen J 2013 Drivers of the dynamics in net primary productivity across ecological zones on the Mongolian Plateau Landscape *Ecol.* 1–15



- Wang S P, Yanfen W, Yonghong L and Zuo-Zhong C 1998 The influence of different stocking rates on herbage regrowth and aboveground net primary production *Acta Agrestia Sinica* **6** 275–81
- Wang Y, Zhou G and Jia B 2008 Modeling SOC and NPP responses of meadow steppe to different grazing intensities in Northeast China *Ecol. Model.* **217** 72–8
- Wesche K and Treiber J 2012 Abiotic and biotic determinants of steppe productivity and performance—a view from Central Asia, Eurasian steppes *Ecological Problems and Livelihoods in a Changing World* (Berlin: Springer) pp 3–43
- Xing X, Xu X, Zhang X, Zhou C, Song M, Shao B and Ouyang H 2010 Simulating net primary production of grasslands in northeastern Asia using MODIS data from 2000 to 2005 *J. Geog. Sci.* **20** 193–204
- Yan L, Zhou G and Zhang F 2013 Effects of different grazing intensities on grassland production in China: a meta-analysis *PLoS ONE* **8** e81466
- Zhang F, Ju W, Shen S, Wang S, Yu G and Han S 2012 Variations of terrestrial net primary productivity in East Asia *Terr. Atmos. Ocean. Sci.* **23** 425–537
- Fangmin Z, Weimin J, Shen S, Wang S, Yu G and Han S 2014 How recent climate change influences water use efficiency in East Asia *Theor. Appl. Climatol.* **116** 359–70
- Zhang N, Zhao Y-S and Yu G-R 2009 Simulated annual carbon fluxes of grassland ecosystems in extremely arid conditions *Ecol. Res.* **24** 185–206
- Zhou G, Zhou X, He Y, Shao J, Hu Z, Liu R, Zhou H and Hosseinibai S 2016 Grazing intensity significantly affects belowground carbon and nitrogen cycling in grassland ecosystems: a meta-analysis *Glob. Change Biol.* **23** 1167–79
- Zhou W, Gang C, Chen Y, Mu S, Sun Z and Li J 2014 Grassland coverage inter-annual variation and its coupling relation with hydrothermal factors in China during 1982–2010 *J. Geog. Sci.* **24** 593–611